



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

QUERY BY PROF. A. HALL.—(1). If  $v$  be the potential function we have the equation given by Laplace

$$\frac{d^2 v}{dx^2} + \frac{d^2 v}{dy^2} + \frac{d^2 v}{dz^2} = 0,$$

which holds for a point outside the attracting body. For a point inside this body we have the equation given by Poisson,

$$\frac{d^2 v}{dx^2} + \frac{d^2 v}{dy^2} + \frac{d^2 v}{dz^2} = -4\pi\rho.$$

What is the value of the right hand side of this equation for a point on the surface of the attracting body? Moigno says, "in this case the expression will have in reality eight distinct values." Statics, p. 460.

(2) Given a hemispherical dome turning about a pin at the top, and having a slit extending from the horizon to the zenith. Can a telescope be placed inside this dome in such a position that every point of the heavens can be seen through the telescope?

QUERY BY T. W. WRIGHT, U. S. LAKE SURV. DETROIT, MICH.—Can a satisfactory proof be given of the statement that "the weight of an observation varies as the square of the probability of its accuracy"? (Monthly Notices R. A. S., Vol. XIII.)

QUERY BY PROF. WM. WOOLSEY JOHNSON, ANNAPOLIS, MD.—"It is stated in Brande and Cox' Dictionary of Sc. and Lit., with respect to Fermat's 'lost theorem', viz.: that  $x^n + y^n = z^n$  is insoluble in integers except when  $n=2$ , that no complete demonstration has yet been given. Yet Barlow in his Theory of Numbers gives what professes to be a demonstration of the impossibility of the equivalent equation  $x^n - y^n = z^n$ , p. 164.

What is the fallacy in Barlow's demonstration and by whom was it first exposed?"

---

### PROBLEMS.

---

313. *By Prof. De Volson Wood.*—A piston, weight  $w$ , is dropped into the end of a vertical cylinder filled with air, length  $l$ ; how far will the piston descend, assuming no friction nor escape of air, nor heat from the compressed air?

314. *By Prof. L. G. Barbour, Richmond, Ky.*—A helix is coiled about a right cylinder the radius of whose base is 1. The projection of the evolute of the helix on the plane of the base of the cylinder encloses double the area of that base. Required the angle made with the plane of the base by the tangent to the helix.